arXiv:hep-th/0208226v1 30 Aug 2002

On new gravitational instantons describing creation of brane-worlds

Mariam Bouhmadi–López*, Pedro F. González–Díaz†
and Alexander Zhuk‡

Instituto de Matemáticas y Física Fundamental,
Consejo Superior de Investigaciones Científicas,
C/ Serrano 121, 28006 Madrid, Spain

2002

Abstract

By considering 5–dimensional cosmological models with a bulk filled with a pressureless scalar field; equivalently dust matter, and a negative cosmological constant, we have found a regular instantonic solution which is free from any singularity at the origin of the extra–coordinate. This instanton describes 5–dimensional asymptotically anti de Sitter wormhole, when the bulk has a topology $R \times S^4$. Compactified brane-world instantons which are built up from such instantonic solution describe either a single brane or a string of branes. Their analytical continuation to the pseudo–Riemannian metric can give rise to either 4-dimensional inflating branes or solutions with the same dynamical behaviour for extra–dimension and branes, in addition to multitemporal solutions. Dust brane-world models with arbitrary dimensions ($D \geq 5$) as well as other spatial topologies are also briefly discussed.

PACS number(s): 04.50.+h, 98.80.Hw

1 Introduction

Although higher dimensional cosmological models can be traced back to the first years of gravitation theory, the idea has received a renewed great attention in the last few years, due to the publication of pioneering papers on brane and extra-dimension models which shed light for the solution of fundamental physical problems including the extra-dimension compactification and the hierarchy problem. Extra-dimensions can be compactified, as in the standard Kaluza-Klein theory [1], or not, as was firstly suggested by Akama [2], Rubakov and Shaposhnikov [3] and others [4]. While Akama [2] considered the universe as a four vortex-like object embedded in a 6–dimensional flat space-time, Rubakov and Shaposhnikov proposed a model in (1+N)–dimensional Minkowski space-time ($N \geq 4$) where particles were confined in a potential well, flat on the usual three spatial dimensions and narrow along the extra-dimensions. Gravity was later on included in 5–dimensional manifolds [4] in order to trap particles near a 4–dimensional Lorentzian submanifolds.

More recently, a model was proposed [5] on (4+N)–dimensions with $N \geq 2$, where the extra-dimensions are compact and their size, R, is deduced by imposing that the usual Planck scale, M_{Pl}, is no longer a fundamental scale and Planck scale in (4+N)–dimensions, $M_{Pl_{4+N}}$, is of the order of the weak scale M_{EW}, $M_{Pl}^2 = M_{Pl_{4+N}}^{2+N} R^N$. This model can solve the hierarchy problem. In this framework the gravitons can propagate in the extra–dimensions while the standard model fields are confined to a 4–dimensional submanifolds of thickness M_{EW}^{-1} in the extra–dimensions.

Randall and Sundrum suggested a new approach to solve the hierarchy problem [6] by including just one extra compact dimension. In their first model [6], inspired by string theory [7], they considered 5–dimensional anti de Sitter (AdS) bulk with two branes with opposite tension; our universe is then placed on the brane with negative tension where standard model particles are localized. In a second model [8], these authors placed the universe in the brane with positive tension, in a non-compact background. In this framework, it is possible to reproduce 4–dimensional general relativity even if the extra–dimension is non-compact [8], due to the existence of a massless gravitational bound state of Kaluza-Klein (KK) modes which is the graviton of 4–dimensional

*e-mail: mbouhmadi@imaff.cfmac.csic.es
†e-mail: p.gonzalezdiaz@imaff.cfmac.csic.es
‡e-mail: ai_zhuk@imaff.cfmac.csic.es
on leave from: Department of Physics, University of Odessa, 2 Dvoryanskaya St., Odessa 65100, Ukraine

world. While for the non-compact case, KK spectrum is continuous without a gap, for the compact case the KK excitations are quantized.

Branes in the Randall-Sundrum models are 4–dimensional flat space-time and consequently, at least in principle, they can not describe any inflationary universe. Nevertheless, as it was pointing out by Garriga and Sasaki (GS) [9] it is still possible to construct an inflating brane, whose geometry corresponds to a 4–dimensional de Sitter space, surrounded by a 5-dimensional AdS. The Euclidean version of this solution can be used for the description of the creation of the universe from nothing. The GS model has a single normalized gravitational bound state which corresponds to the massless graviton and separates by a gap from the massive KK modes.

Multi-brane-world models also exhibit 4–dimensional gravity localization at the branes. In particular, a compact brane-world model consisting of three flat branes, two with positive tensions and an intermediate one with negative tension, embedded in 5-dimensional AdS space-time has been also considered [10]. In this case, the universe is placed in one of the positive tension branes. In addition, intersecting brane configurations [11], crystal brane-world [12] and brane periodic configuration [13] have been investigated as well. The massless gravitational bound state exists in all these models.

However, most of the above mentioned brane-world models have the shortcoming of either not including creation of inflating branes or having a singular character for brane instantons. The main aim of the present paper is to propose a set of singularity free 5–dimensional models which are able to produce inflating brane-worlds. This can be achieved by considering matter in the bulk which we model by a scalar field. As we shall see, such models should also induce localization of 4–dimensional gravity at the branes.

The paper is organized as follows. In the next section we derive a master equation (2.8) which describes the behaviour of the scale factor of D–dimensional instantonic (after an analytic continuation to the Euclidean region) solutions for a bulk filled with a scalar field, which is subjected to a perfect fluid state equation, and a cosmological constant, Λ_D. For $D = 5$, $\Lambda_D < 0$ and spherical 4-dimensional sections, we obtain asymptotically AdS wormhole when the bulk is filled with dust. Using this wormhole solution, in section 3, we construct compact and non-compact brane-world instantons. In section 4, we describe the brane-world birth from nothing (the Euclidean solution) performing an analytical continuation of the instantonic solution. In section 5, we analyze the behaviour of the massless gravitational KK mode for the models constructed in the previous section. Finally, in section 6, we summarize our results and discuss other dust brane-world models with arbitrary dimensions ($D \geq 5$) as well as other spatial topologies.

2 Multidimensional perfect fluid cosmology

Let us start our investigation of a multidimensional model with a cosmological constant, Λ_D, and minimal scalar field, φ, by writing down the action[1]:

$$\begin{aligned} S &= \frac{1}{2\kappa_D^2}\int d^{\,D}X\sqrt{|g^{(D)}|}\left\{R[g^{(D)}] - 2\Lambda_D\right\} \\ &+ \int d^{\,D}X\sqrt{|g^{(D)}|}\left\{-\frac{1}{2}g^{(D)MN}\partial_M\varphi\partial_N\varphi - V(\varphi)\right\} + S_{YGH}\,, \end{aligned} \tag{2.1}$$

where κ_D^2 is D–dimensional gravitational constant and S_{YGH} the standard York-Gibbons-Hawking boundary term [14]. The $(D = 1 + d)$–dimensional metric $g^{(D)}$ is taken to have the form

$$g^{(D)} = g^{(D)}_{MN}dX^M \otimes dX^N = -e^{2\gamma(\tau)}d\tau \otimes d\tau + e^{2\beta(\tau)}g^{(d)}_{\mu\nu}dx^\mu \otimes dx^\nu\,, \tag{2.2}$$

with $g^{(d)}$ the metric of d-dimensional Einstein space: $R[g^{(d)}] = \lambda d \equiv R_d$, and in the case of constant curvature space parameter λ is normalized as $\lambda = k(d-1)$, with $k = \pm 1, 0$.

Consistent with the metric ansatz (2.2), we suppose that the scalar field is also homogeneous: $\varphi = \varphi(\tau)$. For such a scalar field energy density and pressure are defined as:

$$\begin{aligned} T^0_0 &= -\tfrac{1}{2}e^{-2\gamma}\dot{\varphi}^2 - V(\varphi) \equiv -\rho\,, \\ T^\mu_\mu &= \tfrac{1}{2}e^{-2\gamma}\dot{\varphi}^2 - V(\varphi) \equiv P\,, \qquad \mu = 1,\ldots,d\,, \end{aligned} \tag{2.3}$$

and supposed to satisfy the state equation:

$$P = (\alpha - 1)\rho\,. \tag{2.4}$$

The conservation equation $T^M_{N;M} = 0$ implies $\rho = Aa^{-4\alpha}$ where A is an arbitrary constant and $a(\tau) := \exp[\beta(\tau)]$ is a scale factor. This results in a inverse power-law for the scalar field potencial V in term of the scale factor $a = \exp[\beta(\tau)]$; i.e. $V = (1 - \frac{\alpha}{2})A\,a^{-d\alpha}$, whenever $\alpha > 0$. Obviously, a^d gives the volume of the Universe at any hypersurface $\tau = \text{const}$ up to a spatial prefactor $V_d = \int d^dx\sqrt{|g^{(d)}|}$.

[1] Although the constant minimal scalar field is really equivalent to a cosmological term, it is convenient to single out an explicit cosmological constant in the action.

From the energodominant condition it is usually supposed that $\rho \geq 0$ and $-\rho \leq P \leq \rho$, where the latter condition (which results in $0 \leq \alpha \leq 2$) means that the speed of sound is less than the speed of light. For the aim of generality we will not restrict ourselves to satisfy these conditions because at least the first of them can be violated on the case of a scalar field . For example, $\alpha = 0$ with $\rho < 0$ describes the case of a negative cosmological constant. Depending on the values acquired by the parameter α, the scalar field can mimic different kinds of matter. However, the state equation Eq. (2.4) constrains strongly the form of the potential. Below, we shall get, as an example, the potential $V(\varphi)$ corresponding to a dust-like state equation ($\alpha = 1$). Additionally, it can be shown [15] that Einstein equations for such scalar fields are equivalent to the Einstein equations for a perfect fluid with action:

$$S = \frac{V_d}{\kappa_D^2} \int d\tau \left[\frac{1}{2} e^{-\gamma+\gamma_0} d(1-d)\dot{\beta}^2 - e^{\gamma-\gamma_0} U \right] \equiv \frac{V_d}{\kappa_D^2} \int d\tau L \,, \tag{2.5}$$

in which we performed integration over spatial coordinates x, $\gamma_0 := d\beta$, overdot denotes differentiation with respect to τ and we have introduced a "potential energy"

$$U = e^{2\gamma_0} \left(-\frac{1}{2} R_d e^{-2\beta} + \Lambda_D + \kappa_D^2 \rho \right) . \tag{2.6}$$

The constraint equation corresponding to the Lagrangian in Eq.(2.5) results in the following master equation:

$$\frac{\partial L}{\partial \gamma} = 0 \quad \Longrightarrow \quad \frac{1}{2} e^{-\gamma+\gamma_0} d(1-d)\dot{\beta}^2 + e^{\gamma-\gamma_0} U = 0 \,, \tag{2.7}$$

which is exactly the 00–component of the Einstein equation. The function γ reflects the freedom for the choice of different time gauges: $\gamma = 0$ is the proper time gauge, $\gamma = \gamma_0$ is the harmonic time gauge [16] and $\gamma = \beta$ is the conformal time gauge. Thus, in the proper time gauge, Eq.(2.7) reads

$$\left(\frac{\dot{a}}{a} \right)^2 + \frac{R_d}{d(d-1)} \frac{1}{a^2} + \frac{2}{d(1-d)} \Lambda_D + \frac{2}{d(1-d)} \kappa_D^2 A \, a^{-\alpha d} = 0 \,. \tag{2.8}$$

We will concentrate mainly on 5–dimensional brane-world models with AdS bulk: $D = 5$, $\Lambda_D = \Lambda_5 \equiv -|\Lambda_5| < 0$. The scalar curvature R_d can be an arbitrary constant but we shall consider the particular case of constant curvature 4-D space $R_d = 12k$, restricting mainly to the positive curvature with $k = +1$. For these parameters Eq.(2.8) becomes

$$(\dot{a})^2 + k + \Lambda a^2 - \bar{A}^2 \, a^{-4\alpha+2} = 0 \,, \tag{2.9}$$

where for simplicity we have introduced the following notation: $|\Lambda_5|/6 \equiv \Lambda$ and $(1/6)\kappa_5^2 A \equiv \bar{A}^2$. Some of the particular values α are of special interest because they correspond to important types of matter. For example, in 5–D space-time[2] $\alpha = 5/4$ describes radiation, $\alpha = 1$ corresponds to dust (0–D objects), $\alpha = 3/4$ represents cosmic strings (1–D objects), $\alpha = 1/2$ describes domain walls (2–D objects), $\alpha = 1/4$ corresponds to hyperdomain walls (3–D objects) and $\alpha = 0$ represents vacuum (which can be in some sense considered as a 4–D object).

Now, as we want to construct 5–D brane-world models, we perform the Wick rotation to Euclidean "time" $r : \tau \to -ir$. Where r is to be considered as an extra spatial coordinate orthogonal to 4–D branes, i.e., now hypersurfaces at $r = const$. Then, in proper "time" gauge, metric (2.2) is

$$g^{(5)} = dr \otimes dr + a^2(r) g^{(4)} \tag{2.10}$$

and the Euclidean version of Eq.(2.9) reads

$$(\dot{a})^2 - k - \Lambda a^2 + \bar{A}^2 \, a^{-4\alpha+2} = 0 \,. \tag{2.11}$$

Solutions of this equation describe 5–D instantons which can be used to construct brane-world models. Obviously, the precise form of the resulting instantons depends on the type of perfect fluid we use, i.e., on the choice of the parameter α. The vacuum case $\alpha = 0$ (which here corresponds to a simple redefinition of the cosmological constant: $\Lambda_D + \kappa_D^2 A \to \Lambda_D$) was considered in paper [9] for positive curvature $k = +1$ with solution

$$a(r) = l \, \sinh(r/l) \,, \tag{2.12}$$

where $l := \sqrt{\Lambda^{-1}}$ is the AdS radius.

In the present paper we investigate the case of a pressureless scalar field, i.e. with a dust-like state equation ($\alpha = 1$), and $k = +1$. For this particular case the solution of Eq.(2.11) reads:

$$a(r) = \frac{l}{\sqrt{2}} \left(\sqrt{b} \, \cosh \frac{2r}{l} - 1 \right)^{1/2} , \qquad -\infty < r < +\infty \,, \tag{2.13}$$

[2]See e.g. [17] and references therein for the discussion of the equations of state for different kinds of cosmic defects in usual 4–D universe. In these references a regime where the number of defects per co-moving volume is a constant is considered.

where $b := 1+4\Lambda\bar{A}^2$. It can be easily seen that this solution is reduced to (2.12) in the limit: $\bar{A}^2 \to 0 \Longrightarrow b \to 1$. Metric (2.10) with solution (2.13) is non-singular and describes a wormhole (the integration constant being taken in such a way that $r = 0$ corresponds to the wormhole throat). Having the solution (2.13), we can restore the dust-like scalar field potential following the procedure described in [15]. It yields (in the Euclidean region):

$$V(\varphi) = \frac{2A}{l^4}\frac{(\sqrt{b}-1)\tanh^2(\bar{\varphi}) - \sqrt{b} - 1}{(\sqrt{b}-1)^2\tanh^2(\bar{\varphi}) + (\sqrt{b}+1)^2}\,, \tag{2.14}$$

where $\bar{\varphi} \equiv (l/2)\sqrt{b-1/A}\varphi$ and runs from $-\mathrm{arctanh}\left[\sqrt{b-1}/(\sqrt{b}-1)\right]$ to $\mathrm{arctanh}\left[\sqrt{b-1}/(\sqrt{b}-1)\right]$.

It is useful to present solution (2.13) in the conformal "time" gauge: $dr = \pm a d\eta$. It can be easily seen that

$$a(\eta) = l\, b^{1/4}\,\mathrm{ds}\,(b^{1/4}|\eta|)\,, \qquad 0 \le b^{1/4}|\eta| \le 2K(m)\,, \tag{2.15}$$

satisfies the corresponding equation

$$\left(\frac{da}{d\eta}\right)^2 - a^2 - \Lambda a^4 + \bar{A}^2 = 0\,. \tag{2.16}$$

Function $\mathrm{ds}\,u \equiv \mathrm{ds}\,(u|m)$ (with $m = (b^{1/2}+1)/2b^{1/2}$) is a Jacobian elliptic function and $K(m)$ is the complete elliptic integral of the first kind [18]. For each of the sign of η, there is an one–one correspondence between solution (2.15) and wormhole (2.13). Indeed, coordinates r and η are connected with each other by :

$$b^{1/4}\,|\eta| = F\,(\,\phi\,|\,m\,) = u \quad \Longleftrightarrow \quad \mathrm{sn}\; b^{1/4}|\eta| = \frac{1}{\cosh r/l}\,, \tag{2.17}$$

where $\phi := \arcsin(1/\cosh(r/l))$, $\mathrm{sn}\,u \equiv \mathrm{sn}\,(u|m)$ is a Jacobian Elliptic function and $F\,(\,\phi\,|\,m\,)$ is the incomplete elliptic integral of the first kind. In the limit $\bar{A}^2 \to 0 \Longrightarrow m \to 1$, solution (2.15) tends to $a(\eta) \to l/\sinh|\eta|$, which, in fact, is solution (2.12) expressed in the conformal "time" gauge.

3 Brane-world instantons

In this section, we use solutions (2.13) and (2.15) to construct brane-world instantons. This can be performed by excising regions with $r > L$ for two identical wormholes (2.13) and gluing the remaining two copies along the two 4-spheres at $r = L$. The obtained instanton can be described by the following piecewise continuous function:

$$a(r) = \begin{cases} \frac{l}{\sqrt{2}}\left[\sqrt{b}\,\cosh\left(\frac{2r}{l}\right) - 1\right]^{1/2} & , \qquad -\infty < r \le L \\ \frac{l}{\sqrt{2}}\left[\sqrt{b}\,\cosh\left(\frac{2(2L-r)}{l}\right) - 1\right]^{1/2} & , \qquad L \le r < +\infty \end{cases} \tag{3.1}$$

This function is continuous but not smooth at the gluing "point" $r = L$. Due to the Lanczos-Israel junction condition, this results in the appearance of a 4–D spherical brane at $r = L$, with a tension:

$$T(r=L) \,=\, \frac{1}{\kappa_5^2}\frac{3}{4}\widehat{K}(L) \,=\, \frac{6}{\kappa_5^2 l}\,\frac{\sinh(L/l)\,\cosh(L/l)}{\sinh^2(L/l) + m_1} \,>\, 0\,, \tag{3.2}$$

where $m_1 = 1 - m$, $\widehat{K}(L) \equiv K(L^+) - K(L^-)$, and $K(L) = -4a^{-1}(da/dr)_{r=L}$ is the trace of the extrinsic curvature for the case of 5–D metric (2.10).

Instantonic solution (3.1) is non-compact: the scale factor $a(r)$ goes to ∞ when $r \to \pm\infty$, and cannot be used for the description of the brane-world birth from "nothing" (from the Euclidean region) because the probability of it in this case is equal to zero. On the other hand, as we shall see below, the bound state of the spin–2 gravitational perturbations (corresponding to the Newtonian gravity on the brane) is proportional to $a^{3/2}$ and is divergent when $a \to \infty$. However, we can compactify this instanton identifying points corresponding to the throats at $r = 0 \leftrightarrow r = 2L$. It changes the topology along the extra dimension from $\mathbb{R}$ to S^1. Now, the range of the variation of r is the interval $[0,\, 2L]$. Since the geometry is smoothly glued at these points, such procedure does not lead to the appearance of new branes[3].

This procedure for the construction of the brane-world instanton can be easily generalized to the case of an arbitrary number of parallel branes by gluing one-brane manifolds at throats and identifying the two final

[3]Instanton (3.1) can also be compactified by additional cutting at distances $\triangle r = L_1 < L$ from the two sides of the brane, gluing then along these two cuts. Geometry is not smoothly matched at this surface which results in a new negative tension brane. However, in our paper we will not consider such a compact instanton because the first one, with the wormhole throat identification, seems to us more elegant and it leaves a possibility for 5–D baby-universe nucleation at the throat.

opposite throats. For example, in the case of n branes, located at the distances $r = L_i\,,\ i = 1, \ldots, n$ from throats, this instanton can be described by the following piecewise continuous function:

$$a(r) \;=\; \sum_{i=1}^{n+1} a_i(r)\theta_i(r)\,, \qquad\qquad 0 \,\le\, r \,\le\, 2\sum_{i=1}^{n} L_i\,, \tag{3.3}$$

$$a_i(r) \;=\; \frac{l}{\sqrt{2}}\left[\sqrt{b}\,\cosh\left(\frac{2(2\sum_{k=1}^{i-1} L_k - r)}{l}\right) - 1\right]^{1/2}\,, \quad L_{i-1} + 2\sum_{k=1}^{i-2} L_k \,\le\, r \,\le\, L_i + 2\sum_{k=1}^{i-1} L_k\,,$$

where $L_0 \equiv L_{n+1} \equiv 0$ and

$$\theta_i(r) = \tilde{\eta}(r - r_{i-1}) - \tilde{\eta}(r - r_i) = \left\{ \begin{array}{lcl} 0 & , & r < r_{i-1} \\ 1 & , & r_{i-1} \le r < r_i \\ 0 & , & r \ge r_i \end{array} \right. \tag{3.4}$$

are piecewise discontinuous functions, with $\tilde{\eta}(r - r_i)$ being the usual step function[4] equal to zero for $r < r_i$ and becoming unity for $r \ge r_i$. We have redefined the coordinate r in such a way to cover the range of variable for our solution by one coordinate chart. In this case the i-th brane has coordinate $r = r_i = L_i + 2\sum_{k=1}^{i-1} L_k\,,\ i = 1, \ldots, n$ and the i-th throat is located at $r = r_{(th)i} = 2\sum_{k=1}^{i-1} L_k\,,\ i = 1, \ldots, n+1$. "Points" $r = r_0 \equiv r_{(th)1} = 0$ and $r = r_{n+1} \equiv r_{(th)n+1} = 2\sum_{i=1}^{n} L_i$ are identified with each other due to S^1-symmetry. Thus, metric (2.10) with scale factor (3.3) describes a 5–D compact instanton with n parallel branes transversal to coordinate r (see figure 1). Each of these branes has a tension given by Eq.(3.2) with the evident substitution $L \to L_i$ for the i-th (i=1,. . . ,n) brane.

The similar procedure for construction of brane-world instanton can be performed in the conformal gauge. For example, one-brane instanton can be obtained from the solution (2.15) if we take two wormholes with $\eta \lessgtr 0$ and $n = 0$, cut them at $|\eta| = \eta_0 < b^{-1/4}K(m)$, excise regions $|\eta| < \eta_0$ and glue them along this cut. The obtained brane-world instanton is:

$$a(\eta) = l\, b^{1/4}\ \mathrm{ds}\left[b^{1/4}(|\eta| + \eta_0)\right]\,, \tag{3.5}$$

where we redefined the coordinate η in such a way ($|\eta|_{old} \to |\eta|_{new} + \eta_0$) to cover this instanton by one coordinate chart. Here, $0 \le b^{1/4}|\eta| \le 2K(m) - b^{1/4}\eta_0$ and $0 \le b^{1/4}|\eta| \le K(m) - b^{1/4}\eta_0$ (with the identification $-b^{-1/4}K(m) + \eta_0 \leftrightarrow b^{-1/4}K(m) - \eta_0$) correspond respectively to non-compact and compact instantons with brane at $\eta = 0$ and tension

$$T(\eta = 0) \;=\; \frac{6}{\kappa_5^2 l}\,\frac{\sqrt{1 - \mathrm{sn}^2\,(b^{1/4}\eta_0)}}{1 - m\,\mathrm{sn}^2\,(b^{1/4}\eta_0)} \;>\; 0\,, \tag{3.6}$$

which coincides with Eq.(3.2) if $\eta = \eta_0$ and $r = L$ are connected by formula (2.17).

4 Brane-world birth from "nothing"

In this section we investigate the possibility for the creation of brane worlds from "nothing". To be more precise, we interpret the brane-world instantons described in section 3 as a semiclassical paths for the quantum tunneling from the Euclidean region ("nothing").

Euclidean metric (2.10) was obtained from the Lorentzian one by the Wick rotation $\tau \to -ir$. For the compact solutions of section 3 this Euclidean metric describes the compact 5–D brane-world instantons. It is clear that back Wick rotation $r \to i\tau$ will destroy these branes leaving, instead of the brane, a solution corresponding to 5–D baby universe branching off from the wormhole throats at $r = 0$. This baby universe represents 5–D FRW-like universe filled with dust and a negative cosmological constant. It has a maximum at time corresponding to the wormhole throat and bounces from a minimum occurring when the scale factor equals to zero (usual FRW-type singularity). However, there is a possibility for the analytical continuation to the Lorentzian space-time which preserves branes. As it was pointed out in paper [9], it can be done for the case when 4–D positive curvature metric $g^{(4)}$ in Eq.(2.10) is the metric of a 4–sphere:

$$ds_E^2 = dr^2 + a^2(r)(d\chi^2 + \sin^2\chi\, d\Omega_{(3)}^2)\,. \tag{4.1}$$

Then, the analytic continuation of the azimuthal coordinate χ:

$$\chi \longrightarrow iHt + \frac{\pi}{2}\,, \tag{4.2}$$

[4] We put a tilde above it to distinguish from the conformal time η. Obviously, $\theta_i^p = \theta_i\ ,\ p > 0\ ;\quad \theta_i\,\theta_j = 0\ ,\ i \ne j \implies a^p = \sum_{i=1}^{n} a_i^p\theta_i\ ,\ \forall p$ and $\theta_i' = \delta(r - r_{i-1}) - \delta(r - r_i)$.

results in a Lorentzian metric with evolving branes:

$$ds_L^2 = dr^2 + H^2 a^2(r)(-dt^2 + \frac{1}{H^2}\cosh^2 Ht\, d\Omega_{(3)}^2)\,. \tag{4.3}$$

Here, parameter H is chosen in such a way that t describes the proper time on the brane. For example, in the one-brane-case (3.1):

$$Ha(r)|_{r=L} = 1 \quad \Longrightarrow \quad H = \frac{1}{a(L)} = \frac{\sqrt{2}}{l}\left(\sqrt{b}\,\cosh\frac{2L}{l} - 1\right)^{-1/2}\,. \tag{4.4}$$

The compact version of solution (3.1) describes a 4–D spherical brane of radius $a(L) = H^{-1}$ enclosing 5–D space with frozen dust (equivalently, frozen pressureless scalar field) and negative cosmological constant (AdS bulk). The south pole of the 4–D sphere with the azimuth coordinate $\chi = 0$ can be treated as "nothing". The birth of the brane-world takes place at the time $t = 0$ which corresponds in the Euclidean region to $\chi = \pi/2$ (the equator of the 4–D sphere) where the sections $\chi = \text{const}$ of the 4–D sphere reach their maximum with radius H^{-1}. After birth, the brane represents an evolving 3–D sphere with initial radius H^{-1}. More precisely, it is an inflating de Sitter space with the Hubble constant H (see figure 2).

In the case of n branes described by expression (3.3), we can introduce for each of the coordinate patches between the throats $r_{(th)i} \le r \le r_{(th)i+1}$ the following transformation:

$$\chi \longrightarrow iH_i t_i + \frac{\pi}{2}\,, \tag{4.5}$$

where

$$H_i = \frac{1}{a(r_i)} \tag{4.6}$$

and $r_i \in [r_{(th)i}, r_{(th)i+1}]$ is the position of the i-th brane ($i = 1, \ldots, n$). After this continuation, the Lorentzian metric is:

$$ds_L^2 = dr^2 + H_i^2 a^2(r)(-dt_i^2 + \frac{1}{H_i^2}\cosh^2 H_i t_i\, d\Omega_{(3)}^2)\,, \tag{4.7}$$

where coordinates t_i are glued to each other at the throats:

$$t_i = \frac{a_i}{a_j} t_j\,. \tag{4.8}$$

Thus, each of n branes at $r = r_i$ represents a de Sitter space with its own proper time t_i and own Hubble constant H_i.

To conclude this section, we would like shortly comment some alternative possibilities of the analytic continuation. As we have noted before, the analytical continuation $r \to i\tau$ leads to the creation of a 5–D baby universe with scale factor,

$$a(\tau) = \frac{l}{\sqrt{2}}\left(\sqrt{b}\cos\frac{2\tau}{l} - 1\right)^{1/2}\,, \quad 0 \le \tau \le \frac{l}{2}\arccos\frac{1}{\sqrt{b}}\,. \tag{4.9}$$

From this model we can actually construct a spherical 3–D brane by using a similar procedure to that we have described in section 3; that is excising regions with $\chi > L < \pi$ for two identical 4-sphere and gluing the remaining two copies along 3–spheres $\chi = L$. The brane-world model in this case is described by the metric:

$$ds_L^2 = -d\tau^2 + a^2(\tau)(d\chi^2 + a^2(\chi)\, d\Omega_{(3)}^2)\,, \tag{4.10}$$

where the scale factor $a(\tau)$ is defined by Eq.(4.9) and the scale factor $a(\chi)$ reads

$$a(\chi) = \begin{cases} \sin\chi & , \quad 0 \le \chi \le L < \pi \\ \sin(2L - \chi) & , \quad L \le \chi \le 2L \end{cases} \tag{4.11}$$

In this model, both the additional space as well as 3–D brane have the same dynamical behaviour which is described by Eq.(4.9).

We turn finally to another interesting possibility consisting in the simultaneous analytic continuation in two directions: $\chi \to it + \pi/2$ and $r \to i\tau$. It results in a metric:

$$ds_L^2 = -d\tau^2 + a^2(\tau)(-dt^2 + \cosh^2 t\, d\Omega_{(3)}^2)\,, \tag{4.12}$$

which describes a multidimensional/multitemporal solution of the Einstein equations. Relative to time τ we have the FRW-type universe (4.9) and with respect to time t we obtain the de Sitter universe with the Hubble constant $H = 1$. The number of spatial coordinates here is equal to usual three. In the present paper we shall not investigate in detail solutions (4.10) and (4.12) postponing their study for future work.

5 Volcano and multi-Volcano potentials

In this section we will discuss briefly the localization of gravity at the branes described above. It is well known for metrics of the form (2.2) and (2.10), rewritten in the conformal time gauge, that if tensor metric perturbations $\delta g_{MN}(X) = h_{MN}(\eta, x)$ is taken in the gauge: $h_{\eta\eta} = h_{\eta\mu} = h^{\mu}_{\mu} = h^{\mu\nu}{}_{;\nu} = 0$ (where the semicolon denotes a covariant derivative with respect to d–dimensional metric $g^{(d)}_{\mu\nu}(x)$) and is normalized and factorized as $h_{\mu\nu}(\eta, x) = a^{-(d-5)/2}(\eta)\psi(\eta)\hat{h}_{\mu\nu}(x)$, then the radial profile of the perturbations satisfies a Schrödinger-like equation [8]

$$\left(-\frac{d^2}{d\eta^2} + V(\eta)\right)\psi(\eta) = M^2\psi(\eta)\,, \tag{5.1}$$

in which $V(\eta)$ is the so-called "volcano" potential

$$V(\eta) = \left(\frac{d^2}{d\eta^2}a^{\frac{d-1}{2}}\right)\Big/\, a^{\frac{d-1}{2}}\,, \tag{5.2}$$

and M defines a mass spectrum of spin–2 perturbations $\hat{h}_{\mu\nu}(x)$ moving in the d–dimensional background metric $g^{(d)}_{\mu\nu}(x)$. If the spectrum of mass starts from $M = 0$, then a state corresponding to this value of M describes usual 4–dimensional (for $d = 4$) massless gravitons responsible for the standard Newtonian gravity.

It can be seen that for $M = 0$ the trivial solution of Eq.(5.1) is

$$\psi \propto a^{\frac{d-1}{2}}\,. \tag{5.3}$$

Thus, if the scale factor a has maxima at branes, this zero-mode state is localized at the branes. On the other hand, if the shape of the volcano potential is such that it suppresses the Kaluza–Klein modes with $M > 0$ at the branes by the wings of the potential, then the massless graviton gives the main contribution to gravity on the brane and effectively we have the Newtonian gravitation here. Moreover, to have a well defined effective gravitation on branes, it is desirable to have a mass spectrum where the massless graviton is well separated by a mass gap from other massive Kaluza–Klein modes [13]. This is precisely the general situation that we have in the present paper for compact models. Indeed, in the case of the compact n-brane model described by (3.3) the radial profile reads:

$$\psi \propto a^{3/2}(r) = \sum_{i=1}^{n+1} a_i^{3/2}(r)\,\theta_i(r)\,, \tag{5.4}$$

where we used the properties of the θ–function (see footnote 4). This formula shows that the massless graviton (zero-mode) is localized on each of the branes as they approaches their local maximum at the branes.

The volcano potential (5.2) for this model can be written as follows

$$\begin{aligned} V &= a^{-3/2}\frac{d^2}{d\eta^2}a^{3/2} = a^{-3/2}\left(a\frac{da}{dr}\frac{d}{dr}a^{3/2} + a^2\frac{d^2}{dr^2}a^{3/2}\right) \\ &= \sum_{i=1}^{n+1}\overline{V}_i(r)\theta_i(r) - \frac{16}{3}\kappa_5^2\sum_{i=1}^{n} a_i^2 T(r)\delta(r - r_i)\,, \end{aligned} \tag{5.5}$$

where

$$\overline{V}_i(r) = \frac{15}{4}\sqrt{b}\cosh^2(r - 2\sum_{k=1}^{i-1}L_k)/l\ \ \frac{\sinh^2(r - 2\sum_{k=1}^{i-1}L_k)/l + (4/5)m_1}{\sinh^2(r - 2\sum_{k=1}^{i-1}L_k)/l + m_1}\ -\ \frac{3}{2}\sqrt{b}\,. \tag{5.6}$$

Here, we used the expressions (3.3) and (3.4) for the functions a_i and θ_i and the tension $T(r_i)$ has the form of Eq.(3.2) with the replacement $L \to L_i\,,\ i = 1, \ldots, n$. It can be seen that, at the throats, the potential (5.5) reaches its minima $V(r)|_{r_{(th)i}}\,,\ i = 1, \ldots, n$:

$$V(r)|_{r_{(th)i}}\ \equiv V_{th}\ =\ \begin{cases} (3/2)\sqrt{b} & ,\quad b \neq 1 \\ 9/4 & ,\quad b = 1 \end{cases} \tag{5.7}$$

and, at the branes, achieves its local maxima:

$$V_{(br)i} = \frac{15}{4}\sqrt{b}\cosh^2(L_i/l)\frac{\sinh^2(L_i/l) + (4/5)m_1}{\sinh^2(L_i/l) + m_1}\ -\ \frac{3}{2}\sqrt{b}\,,\quad i = 1, \ldots, n\,, \tag{5.8}$$

which tend to $(15/4)\cosh^2(L_i/l) - 3/2$ in the limit $b \to 1$, corresponding to the vacuum case (2.12). Two different values of the minimum expressed in Eq. (5.7) result from the two different asymptotic behaviours of the function $\mathrm{cd}\,(u|m) \to 0/\sqrt{1-m}$ when $u \to K(m)$. Hence, this fraction equals to 0 if $m \neq 1$ and goes to 1

when $m \to 1$. The second value of Eq. (5.7) $V_1 = 9/4$ was obtained in [9] for model (2.12). Thus, Kaluza–Klein modes starts at

$$M = \sqrt{V_{th}}\,, \tag{5.9}$$

and this mass gap separates the zero-mode from other massive Kaluza–Klein modes[5]. It is obvious that the spectrum of the Kaluza–Klein modes is discrete for the compact models.

Thus, in the n-brane case, potential (5.5) has the form of a string (closed string for compact models) of volcano potentials (see figure 3) and can be named as "multi-Volcano potential". Obviously, this form of the potential entails suppression of the massive Kaluza-Klein modes at the branes.

To conclude this section, we would like to remark an interesting possibility arising from the special form of the multi-Volcano potential with periodic structure. In this case we obtain a band structure for the mass spectrum of the Kaluza-Klein modes and only masses from these bands are allowed to exist (see e.g. [13]). This would give rise to separation between the zero-mode and the massive Kaluza–Klein modes even larger than those induced by the mass gap (5.9).

6 Conclusions

In this paper we have considered in some detail 5–dimensional cosmological models corresponding to the case of a bulk filled with a pressureless scalar field, or equivalently by dust, and a cosmological constant. Special attention has been given to the case of a negative cosmological constant in a bulk with the spatial geometry of a four-sphere. For this model, we have found an asymptotically AdS wormhole instantonic solution. Brane-world instantons with a single 4–D spherical brane as well as with a string of such concentric branes can then be built up by using a cutting and gluing procedure. We have been able to obtain regular solutions which are free from any singularities at the origin of extra coordinates, and can be compactified so that the asymptotic divergences of the scale factor are prevented. Zero-mode massless gravitons are shown to be localized on these 4–D branes, so allowing such branes to nest Newtonian gravity. Analytical continuation from the brane instantonic metric to the Lorentzian regime leads to de Sitter 4–dimensional inflating branes. After birth of the inflating brane world from "nothing", the perfect fluid (dust) remains frozen: it is contained in the bulk but not on the brane. Here, inflation has pure geometrical origin: the Hubble constant of the inflating brane world is defined by the inverse radius of the Euclidean 4–D spherical brane.

Other brane-world models with similar proprieties, as the ones discussed in the sections 3 and 4, can be obtained following the procedure explained in the present paper. For example, inflating branes can be constructed when the 5–D space has positive or zero cosmological constant, whenever the topology of the instanton is $\mathbb{R}\times S^4$. Moreover, using these instantons, we can get a class of brane-world models characterized by a common dynamical behaviour for the extra-dimension and brane. On the other hand, brane-world models with static flat brane or with a number of such parallel branes can be described considering an instanton with negative cosmological constant and flat 4–D spatial topology ($k = 0$).

Additionally, our model can be generalized for an arbitrary number of dimensions. This can be performed by considering the model (2.2) with topology $\mathbb{R}\times M^d$. Here, the d-dimensional manifold M^d undergoes a topological splitting into n Einstein spaces: $M^d = \prod_{i=1}^n M_i^{d_i}$, $g^{(d)} = \sum_{i=1}^n g_{(i)}^{(d_i)}$, $R_{\mu\nu}[g_{(i)}^{(d_i)}] = \lambda_i g_{(i)\mu\nu}^{(d_i)}$, $d = \sum_{i=1}^n d_i$, with $\lambda_i > 0$. Now, using the conformal transformation $g_{(i)}^{(d_i)} \to (\lambda_i/\lambda) g_{(i)}^{(d_i)}$, it can be easily shown [20] that the constituent manifold M^d is also the Einstein space: $R_{\mu\nu}[g^{(d)}] = \lambda g_{\mu\nu}^{(d)}$ and the scale factor $a(\tau)$ satisfies Eq. (2.8). Thus, the brane-world instantons, as well as the birth of the brane-worlds from them, can be constructed for this $(d+1)$-dimensional model in complete analogy to the sections 3 and 4. For example, in the particular case $M^d = S^4 \times \prod_{i=1}^n S^{d_i}$, the analytic continuation (4.2) with respect to the coordinate χ of 4–sphere S^4 results in the following Lorentzian metric:

$$ds_L^2 = dr^2 + H^2 a^2(r)(-dt^2 + \frac{1}{H^2}\cosh^2 Ht\; d\Omega_{(3)}^2 + \frac{r_1^2}{H^2}\; d\Omega_{(d_1)}^2 + \ldots + \frac{r_n^2}{H^2}\; d\Omega_{(d_n)}^2)\,, \tag{6.1}$$

where $r_i\,,\ i = 1, \ldots, n$ are the radii of d_i-spheres. We arrive at the brane-world model with inflating 4–D part of the brane, plus frozen, compactified and unobservable (for $r_i \lesssim 10^{-17} cm$) $(d-4)$ dimensions on it. This scenario, with a brane of codimension one, is of interest because:

1. it demonstrates the very interesting possibility for the unification of new brane-world scenarios with the standard Kaluza–Klein approach.
2. it gives a possibility for the localization of the gauge fields on the brane [21], [22].

Of course, some important aspects of this research need further investigations including the study of the transition from inflationary branes into a matter-dominated brane universe, the physical consequences from the above mentioned brane-world model with dynamical equivalence between extra dimension and branes, as well as the meaning of the 5-dimensional space-times described by a metric with two timelike dimensions.

[5]It is clear, that to an observer on a i-th brane at $r = r_i$, the physical metric is the induced metric on this brane: $g_{(ph)\mu\nu}^{(4)} = a_i^2(r_i) g_{\mu\nu}^{(4)}$. It results in an appropriate rescaling of effective physical values of, for example, effective 4–D cosmological and gravitation constants on the brane (see [19]). Obviously, physical Kaluza–Klein masses for this observer are similarly rescaled, $M \to m_{(ph)} = M/a_i(r_i)$.

Acknowledgments

A.Z. thanks Instituto de Matemáticas y Física Fundamental, CSIC, for kind hospitality during preparation of this paper. A.Z. acknowledges support by Spanish Ministry of Education, Culture and Sport (the programme for Sabbatical Stay in Spain) and the programme SCOPES (Scientific co-operation between Eastern Europe and Switzerland) of the Swiss National Science Foundation, project No. 7SUPJ062239. M.B.L. is supported by a grant of the Spanish Ministry of Science and Technology. This investigation was supported by the GICYT under Research Project No. PB97-1218.

References

[1] Kaluza Th. Zum Unitätsproblem der Physik, Sitzungsber. d. Preuss. Akad. d. Wiss., 1921, 966.
Klein O. Quantentheorie and funfdimensinale Relativitatstheorie, Zeitschrift für Physik., 1926, V. 37, 895-906.

[2] K. Akama, *Pregeometry*, (Lectures Notes in Physics, Gauge, Theory and Gravitation, Proc. Int. Symp. *Gauge Theory and Gravitation*, Nara, Japan, August 20-24, 1982), Springer-Verlag, 1983, Ed. K. Kikkawa, N. Nakanishi and Nariai, P. 267 - 271.

[3] V. A. Rubakov and M.E. Shaposhnikov, Phys. Lett. B125, (1983), 136 - 138.

[4] M. Visser, Phys. Lett. B156, (1985), 22 - 25;
E.J. Squires, Phys. Lett. B167, (1986), 286 - 288;
M.D. Maia and V. Silveira, Phys. Rev D48, (1993), 954 - 957.

[5] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B429, (1998), 263 - 272, hep-ph/9803315.

[6] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, (1999), 3370 - 3373, hep-ph/9905221.

[7] P. Hořava and E. Witten, Nucl. Phys. B460, (1996), 506 - 524, hep-th/9510209;
E. Witten, Nucl. Phys. B471, (1996), 135 - 158, hep-th/9602070;
P. Hořava and E. Witten, Nucl. Phys. B475, (1996), 94 - 114, hep-th/9603142.

[8] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, (1999), 4690, hep-th/9906064.

[9] J. Garriga and M. Sasaki, Phys. Rev. D62, (2000), 043523, hep-th/9912118.

[10] I.I. Kogan, S. Mouslopoulos, A. Papazoglou, G.G. Ross and J. Santiago, Nucl. Phys. B584, (2000), 313 - 328, hep-ph/9912552.

[11] N. Arkani-Hamed, S. Dimopoulos, G. Dvali and N. Kaloper, Phys. Rev. Lett. 84, (2000), 586 - 589, hep-th/9907209.

[12] N. Kaloper, Phys. Lett. B474, (2000), 268 - 281, hep-th/9912125.

[13] S. Nam, JHEP 0004, (2000), 002, hep-th/9911237.

[14] J.W. York, Phys. Rev. Lett. 28, (1972), 1082 - 1085;
G.W. Gibbons and S.W. Hawking, Phys. Rev. D15, (1977), 2752 - 2756.

[15] A. Zhuk, Class. Quant. Grav. 13, (1996), 2163 - 2178.

[16] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, Nuovo Cim. B104, (1989), 575.

[17] R.A. Battye, M.Bucher and D. Spergel *Domain wall dominated universe*, asrto-ph/9908047.
M. Bucher and D.N. Spergel, Phys. Rev. D60, (1999), 043505, astro-ph/ 9812022.

[18] M.Abramowitz and I.A. Stegun *Handbook of Matematical Functions* , Dover Publ. Inc., New York, 1972.

[19] M. Bouhmadi–López and A. Zhuk, Phys.Rev. D65, (2002), 044009, hep-th/0107227.

[20] A. Zhuk, Grav.Cosmol. 3, (1997), 24-28, gr-qc/9701037.

[21] S.L. Dubovsky, V.A. Rubakov and P.G. Tinyakov, JHEP 0008, (2000), 041, hep-ph/0007179.

[22] V.A. Rubakov Phys.Usp. 44, (2001), 871-893; Usp.Fiz.Nauk 171, (2001), 913-938, hep-ph/0104152.



Figure 1: Compact n-brane-world instanton (3.3) in the case $n = 3$. Each point in the figure represents a 3–D sphere. Lines with fixed coordinates $r = \text{const} \equiv r_0$ correspond to 4–D spheres with radii $a(r_0)$. The branes are 4–D concentric spheres surrounding the 5–D bulk.



Figure 2: Birth of the one-brane-world ($n = 1$) from "nothing". The creation takes place at the time $t = 0$ and the 5–D Lorentzian metric is described by Eq.(4.3). The brane after birth represents an inflating 3–D sphere with initial radius H^{-1} that is de Sitter space-time with Hubble constant H.



Figure 3: Multi-Volcano potential (5.5) for the compact 5–brane-world instanton when $n = 5$. Each of the "volcanoes" has a local maximum $V_{(br)i}$ localized on the branes. Identical local minima V_{th} correspond to the wormhole throat positions $r_{(th)i}$.